LASSO* – Science Requirements

*LES ARM Symbiotic Simulation and Observation (LASSO) workflow



Andy Vogelmann¹, William I Gustafson Jr²
Zhijin Li^{3,4}, Xiaoping Cheng³,
Satoshi Endo¹, Tami Toto¹, and Heng Xiao²



¹Brookhaven National Laboratory ²Pacific Northwest National Laboratory ³University of California Los Angeles ⁴NASA Jet Propulsion Laboratory

And TONS of people from the rest of ARM!

LASSO Webpage: http://www.arm.gov/science/themes/lasso

LASSO e-mail list sign up: http://eepurl.com/bCS8s5

LASSO Overview

Goal of routine LES modeling at the ARM sites

- Add value to ARM observations by using LES modeling to provide context and a *self-consistent representation* of the atmosphere that connects processes and facilitates improved understanding.
- Produce a library of simulations to support the statistical study of atmospheric processes and support the improvement of the parameterizations of these processes in climate models. Also serve as a proxy for the atmosphere to develop remote sensing retrievals.


The plan

- Starts at the Southern Great Plains megasite in 2017
- Initial implementation targets shallow clouds
- Later expands to other phenomena and ARM sites

ARM observations integral to the effort

- Model initialization and forcings, including a data assimilation effort
- Evaluation of simulations
- Generation of model-observation "data cubes" for analysis by the community

LASSO Workflow: 2-Year Pilot Phase




Major focus on getting something that works in place and improve it with time

- Implement things that already work
- > Do not have time for experimenting with new/untested things

LASSO Data Priorities

- Model forcing and initialization
- Model evaluation



Prioritization of Radar-Related Observables

Value

1 = Critical

2 = Important

3 = Nice to have

Continuity

1 = Always

2 = Event conditionality

3 = "Irregular" conditionality

Keep in mind: large-eddy simulations are statistical in nature

- Biases are baaaaaaad
- RMS consistent with modeling framework, given adequate sampling



Target Properties and Specifications (1)


	Value	Continuity	Desired Specifications	Comments			
Model Forcing & Initialization							
Horizontal Winds	1	1	Accuracy of 1 m/s and 3° Vertical resolution: 30-40 m in BL 100 m above	Must be all-weather → RWP ~10-15 min averages Also gets virtual temperature to 0.5 K			
In-Cloud Dynamics (instrument/method simulator)							
Vertical velocity	1	2	≤ 20%	Range-gate resolved Turbulence and entrainment value not "1" (yet) because of potential issues matching with model computations			
Mass flux	1	2					
Turbulence	1-2	2-3	≤ 50%				
Entrainment	1-2	2-3	≤ 50%				

Target Properties and Specifications (2)

	Value	Continuity	Desired Specifications	Comments			
Cloud Boundaries (radar simulator)							
Cloud thickness/ Cloud-top height	1-2	1-2	≤ 50-100 m (RACORO clouds ~200-300 m thick)	Value reduced due to model resolution limitation PDFs of 5-min "snapshots" aggregated over 15-20 min Unknown effects of CTH radar detection and insect contamination			
Cloud volume boundaries (3-D)	1-2	2	≤ 100 m (Limited by horizontal model resolution)	Value reduced due to model resolution limitation Cloud-base area needed PDF every 100 m in the vertical Core-center height			
Cloud Microphysics and Precipitation (radar simulator)							
LWC	1	2	Whatever we can get (!), preferably ~0.1 gm ⁻³ (median for RACORO clouds)	Range-gate resolved			
Drizzle	2	2	Projected area of occurrence ~20%	Value limited by drizzle frequency at			
	2	3	Drizzle rate at cloud base ~100%	SGP (vs. ENA) Area more important than amount			
Droplet number concentration	1	3	MIN[50%, < 250 cm ⁻³], but anything would help (RACORO range was 100-1000 cm ⁻³)	Conditional on LWC detection and probably other factors that would affect continuity			

Synthesis Products

For example, ENCORE (ENsemble ClOud REtrieval method)



Summary

This is a work in progress

We are interested in your feedback!

vogelmann@bnl.gov

William.Gustafson@pnnl.gov

LASSO Webpage: http://www.arm.gov/science/themes/lasso

LASSO e-mail list sign up: http://eepurl.com/bCS8s5

Unused Slides

Highlights of the Pilot Project's 2-Year Timeframe

May 2015 Funding started

May 2016 ARM/ASR Joint Meeting May – Coming out party!

June 2016 Availability of initial simulations for shallow convection

cases during spring-summer 2015 (6-10 cases) with

associated forcing, observational data.


→ Beta users (including ASR call, CMDV call)

January 2017 Simulations available for spring—summer 2016 that

include input data from the new boundary facilities.

May 2017 Pilot phase completed

LASSO ARM Data Infusion



Four main ARM data uses

a. Scene classification

Cloud classification VAP (Sunny & Laura)

b. Forcing generation & Initialization

Atmos state profiles (T, Q, horizontal winds)

SH & LH or soil moisture (Noah LS)

Other surface observations (Mesonet)

Variation of initial conditions

c. Evaluation diagnostics & metrics

Cloud properties

Vertical velocities


Other unassimilated variables

d. Data cube generation

Observations

Observation-comparable model variables

Other non-comparable model fields



Model Forcing Generation

MS-DA needs ARM atmospheric observations

Profiles of T, Q, and horizontal winds

Wind profilers needed for boundary layer winds

- Mixture of precip mode with BL mode effective?
- → January 2017 deadline for delivery of initial simulations using new boundary facility data

